

Clenpiq Bowel Preparation Instructions – Morning Exams

DO NOT follow the instructions on the box.

You are scheduled for a colonoscopy with Dr. _____ on _____.

Your scheduled time is _____, but please arrive at _____.

- Fort Sanders Regional Hospital. Please see patient services center in the main lobby.
- Centers for Advanced Medicine (CAM). Suite 218, on second floor.
- LeConte Medical Center in Sevierville. Go in the front door to registration desk.

Medications

- You may take your heart, blood pressure, seizure, thyroid, and reflux the morning of your procedure.
- **Do NOT take blood thinners including: Coumadin® (warfarin); Plavix®(clopidogrel); Brilinta (Ticagrelor); Ticlid® (ticlopidine hydrochloride); Agrylin® (anagrelide); Xarelto® (Rivaroxaban); Pradaxa® (Dabigatran); Eliquis® (Apixaban); Effient® (Prasugrel); and Aspirin. Stop Medication _____.**

If you are diabetic:

- If you take a pill to lower your sugar, do not take it on the day of your procedure.
- If you are taking regular insulin®, do not take it on the day of your procedure.
- If you are taking long-acting insulin (NPH), take half of your prescribed dose on the morning of the procedure.
- If you are taking any other insulin preparation, such as Lantus®, Humalog® or 70/30 insulin, contact your prescribing physician for instructions.

Seven (7) days before your colonoscopy

- If you experience constipation, begin Miralax 1 to 2 times a day, 7 days prior to your scope.

Five (5) days before your colonoscopy

- Do **NOT** take medicines that stop diarrhea, such as loperamide (Imodium®), Kaopectate® or Pepto-Bismol®.
- Do **NOT** take fiber supplements, such as Metamucil®, Citrucel®, or Perdiem®.
- Do **NOT** take products that contain iron, such as multivitamins, ferrous sulfate, iron supplements, etc.
- Do **NOT** take Vitamin E.

Three (3) days before your colonoscopy: No Fibrous Foods or Drinks **** Please refer to chart labeled CHART 1 ****

One (1) day before your colonoscopy

- **Do NOT eat any solid foods.** Drink at least 8 ounces of clear liquids every hour after waking up. **** Please refer to chart labeled CHART 2****

Part 1 at 3 PM on the evening before your colonoscopy

1. Drink one bottle of CLENPIQ®.
2. Over the next five hours, drink at least 5 cups (8-ounces each) of clear liquid at your own pace.
3. You may continue to drink clear liquids until midnight.

Part 2 at 7 PM on the evening before your colonoscopy:

1. Drink one bottle of CLENPIQ, then drink one cup (8-ounces each) of clear liquid, every 15 minutes for at least five cups.
2. You may continue to drink clear liquids up to 4 hours before your exam.

If you have any questions or need to cancel/reschedule your appointment, please call 865-523-6418.

Chart 1	
Ok To Eat	Avoid These Foods
Creamy Peanut Butter, Butter, Margarine, Mayonnaise, Salad Dressing without nuts, Cheese, Yogurt, Sour Cream, Buttermilk, Cream, Hot Chocolate	Nuts, Seeds, Rice, Whole Grain, Chili, Lentil, Dried Beans, Corn, Pea, Legumes, Lima Beans, Tough Meat with Gristle
Broth, bouillon, Consommé, Strained Soups, Milk, Custard, Plain pudding, Ice Cream, Jell-O, Cookies Without Nuts, Coffee, Tea, Hot Chocolate, Clear Fruit Drinks, Soda, Boost or Ensure without Fiber, Sugar, Salt, Jelly, Honey, Syrup, Lemon Juice	Raw Fruits, Cooked or Canned Fruit with Seeds or Skin, Raisins, Corn, Potatoes WITH skin, Tomatoes, Cucumbers with seeds and peel Cooked cabbage or Brussel Sprouts, Squash , Onions,
Ripe, Peeled Fruits and Vegetables, Canned or Cooked Fruit or Vegetables Without Seeds or Skin, Potatoes or Cucumbers without Seeds or Skin	Coconut, popcorn, Jam, Marmalade, Relishes, Pickles, Olives, Stone-ground mustard
Low Fiber Cereal's, Bread, Crackers.	Red or Purple Dye
Chicken, Turkey, Lamb, Lean Pork, Veal, Fish, Seafood, Eggs, Tofu	Pulp

Chart 2	
Ok To Drink	DO NOT DRINK
Gatorade®, Pedialyte® or Powerade®	Alcohol
Clear broth or bouillon	Milk or non-dairy creamer
Coffee or tea (no milk or non-dairy creamer)	Noodles or vegetables in soup
Carbonated and non-carbonated soft drinks	Juice with pulp
Kool-Aid® or other fruit-flavored drinks	Liquid you cannot see through
Strained Fruit Juices (no pulp)	Red/Purple/Blue Liquids
Jell-O®, Popsicles®, Hard Candy	
Water	

If you have any questions or need to cancel/reschedule your appointment, please call 865-523-6418.